

To study the effect of different chemicals and packing materials on the physiological and organoleptic characteristics of mosambi (*Citrus limetta*) fruit

Dr.R Chithiraichelvan

Faculty, Dept of Horticulture,

Bharatiya Engineering Science & Technology Innovation University, Andhra Pradesh, India.

Abstract

The experiment was conducted “to study the effect of different chemicals and packing materials on the physiological and organoleptic characteristics of Mosambi (*Citrus limetta*) fruit” at the Agricultural laboratory of Sant Baba Bhag Singh University, Jalandhar, Punjab (India) and Punjab agro. Juice Ltd. Hoshiarpur FSSC LIC No 10014063000386 and at Lal Path lab, Nandachor, Punjab (India) during the year 2020. The experiment was laid out in Completely Randomized Design (CRD) with three replications and 10 treatments. After washing and drying, fruits are pre-treated with chemicals viz. 2, 4- dichloro phenoxy acetic acid (2, 4-D) and gibberellic acid (GA3) and packed in various packaging materials viz. high-density polyethylene (HDPE) and other packaging materials, Low-density polyethylene (LDPE) with or without perforations) and sealed with an electronic sealing system after being dehumidified. Fruits samples were analysed for various physiological characteristics and organoleptic characters after 7 days of an interval of storage (0, 7th, 14th, and 21th). The result revealed that among the different treatments T₄ (HDPE perforated) treatment was superior for characters like fruit appearance, individual fruit weight, rind thickness, fruit juice content %, colour, and flavor as compared to other treatments in respect of maintaining morphological and organoleptic parameters. However, least values were recorded in T₀ (Control) for most of the parameters analysed. Therefore, treatment of T₄ (HDPE perforated) may be suggested as effective and economical treatment for improving shelf life.

Keywords: LDPE, HDPE and CRD, chemicals and packing materials, improving shelf life

Introduction

Citrus limetta, is considered to be a variety of Sweet orange and citrus varieties, belongs to the Rutaceae family, and is often referred to Mosambi and Sweet limetta. *Citrus limetta* can be grown locally in tropical or subtropical climates, and fruit production begins to appear on plants that are 5 to 7 years old. It starts to evolve and produces a good yield from 10 to 18 years. Global production of sweet oranges in 2014 was 70.9 million tons, 24% of the total global production was produced by Brazil followed by China and India ^[3]. During 2016- 2017, overall area under cultivation of Mosambi was 209.19 / ha with fruit production of 3497.35 MT and productivity 16.7 MT/ha ^[4]. The role of packaging to increase the shelf life of fruit is very crucial in post-harvest operations of horticultural fruit crops. Each polymer-

sealed fruit is better than a few fruits stored in a plastic bag because sealing each fruit allows for a more efficient environment against respiratory biomass and even each sealed packaging helps in preventing the fruit from decaying infections that are considered secondary agents ^[2]. In view of the above fact, the present study was carried out to investigate the effect of different chemicals and packing materials on the physiological and organoleptic characteristics of Mosambi fruit.

Methods and Materials

The experiment was laid out in Completely Randomized Design in which 10 treatments were implemented with 3 replications after 7 days of interval (0, 7th, 14th, and 21th) for identification of better effect of different chemicals and packaging materials on physiological and organoleptic traits of Mosambi in December 2020. Fruits are pre-treatment with hot water (55 °C) and the fruits were air dried room temperature for one hour after pre-treatment.

The mosambi fruits were wrapped in various packaging materials *viz.* high-density polyethylene (HDPE) and other packaging materials, Low-density polyethylene (LDPE) with or without perforations) and sealed with an electronic sealing system after being dehumidified. Chemically *viz.* 2, 4-dichlorophenoxyacetic acid (2, 4-D), and gibberellic acid (GA3), prepared fruits were also kept at (ambient) room temperature. Fruit samples were analysed for various physical characteristics *viz.* Fruit appearance, Fruit diameter, Individual fruit weight, Rind Thickness and Fruit juice % and organoleptic characteristics *viz.* Colour, Flavour, and Overall Acceptability. Therefore, present investigation was conducted with 10 different treatments *viz.*, T₁ (LDPE), T₂ (LDPE perforated), T₃ (HDPE), T₄ (HDPE perforated), T₅ (Cold storage), T₆ (Treated with 2,4-D 50 ppm (without packaging)), T₇ (Treated with 2,4-D 100 ppm (without packaging)), T₈ (Treated with GA3 50 ppm (without packaging)), T₉ (Treated with GA3 100 ppm (without packaging)), T₀ (Control).

Results and Discussion

The analysis of variance revealed significant differences among the treatments for all the traits attributes under study. Data recorded on the effect of different chemicals and packing materials on physiological and organoleptic parameters of Mosambi (*Citrus limetta*) fruit.

Physiological parameter Fruit Appearance

This study shows that chemicals and packaging materials have a significant impact on Mosambi at the end or end of the 21st storage period. Maximum value of appearance (6.78) was obtained in HDPE perforated (T₄) whereas minimum (5.23) was recorded in control (T₀) shown in Table 1. These findings are supported by Tatlı and Ozguven (1999) ^[10], as well as Tugwell and Chvyl (1996) ^[11].

Individual fruit weight

Chemicals and packaging materials were found to have a significant impact on individual fruit weights as shown in Table 2. The minimum reduction in weight from 0 days to 21 days (153.49 gm) was obtained in HDPE perforated treatment (T₄) whereas, minimum value (124.50 gm) in control (T₀) on 21th day of the storage due to maturation and decrease during respiration and respiration. The current results are supported by Bhullar ^[1], who investigated that chemically treated fruits with 2, 4-D, and high-density polyethylene can be stored.

Fruit diameter

The results of this study showed that the packaging material had a significant effect on fruit diameter, but the minimal reduction in fruit diameter found during cold storage (T₅) showed better results than other packaging materials on the other hand in chemical treatment of fruit treated with 2, 4 -D 100 ppm (T₇) showed minimal diameter reduction compared to other pretreatment chemicals. The maximum diameter reduction was found in the control (T₀), (Table 3).

Rind thickness

In the present study, it was found that chemically treated fruits, cold handling, and packaging materials were of very different sizes, as shown in Table 4. Minimal reduction in skin thickness was noted in T₄. The maximum shrinkage of bark thickness was recorded in the treatment or control (T₀). The reduction in skin thickness of unprocessed fruit is maximal under ambient or ambient temperature conditions, resulting in greater moisture loss and skin thickness shrinkage, a term used by Wills *et al.* ^[12].

Fruit juice %

This study showed that juice weight decreased with increasing storage time and showed a significant effect of different treatments on fruit juices as shown in Table 5. The maximum juice content (51.31%) was determined in perforated high-density polyethylene (T₄) and the minimum juice content (46.21%) in the control (T₀) was recorded. This may be due the property of high-density polyethylene, supplying a barrier in opposition to the water vapor and continues relative humidity which decreases the transpiration rate. The current findings are supported through Kaur ^[5] and Singh ^[7] in Kinnow Mandarin.

Organoleptic parameter Colour

The records presented in the Table 6 is indicating that the packaging material used to be appropriate for preserving the fruit colour; fruits sealed in excessive density polyethylene with perforation (T₄) confirmed less color modifications with slower rate. Whereas, the maximum rapid and quick colour

changes regarded in control (T_0). The current findings are supported through the end result obtained by means of the Siddiqui ^[8] and Mandhyan ^[6].

Flavor

The different packaging treatments affected the flavor at the last of storage, large value recorded in packaging material was in high-density polyethylene with perforation (T_4) (5.48), and minimum value was recorded in control (T_0) shown in Table 7. The present findings are supported by Sonkar and Ladaniya ^[9].

Overall acceptability

In this experiment, various treatments of packaging, cold treatments, and chemical treatments significantly affects the fruits overall acceptability shown in Table 8. The maximum value (5.41) of acceptability recorded in cold treatment (T_7) and this may be due to the low temperature, which limits the biochemical activity, reduces the respiration rate of the fruit, causing delays in ripening and the fruit can be stored for days. Whereas, minimum value (2.12) was recorded in control (T_0).

Table 1: Effect of different treatments on appearance of mosambi fruits

Treatments	7 Days	14 days	21 days
T ₁	7.58±0.34	6.94±0.36	6.12±0.65
T ₂	7.49±0.54	6.82±0.54	6.33±0.51
T ₃	7.67±0.23	6.86±0.50	6.48±0.50
T ₄	7.89±0.67	7.10±0.98	6.78±0.65
T ₅	7.41±0.54	6.42±0.56	6.02±0.65
T ₆	7.30±0.23	6.52±0.87	6.24±0.54
T ₇	7.50±0.54	6.98±0.12	6.39±0.31
T ₈	7.42±0.87	6.61±0.37	6.04±0.87
T ₉	7.39±0.76	6.56±0.41	6.12±0.56
T ₀	7.10±0.98	6.18±0.23	5.23±0.98

Table 2: Effect of different treatments on the individual fruit weight (gm)

Treatmen ts	0 Days	7 Days	14 days	21 days
T ₁	172.67±3. 23	166.45±2. 56	158.29±2. 58	151.92±2. 55
T ₂	171.36±2. 98	165.26±3. 65	156.66±1. 59	151.03±3. 79
T ₃	169.22±4. 33	161.45±2. 98	153.87±3. 19	147.65±4. 11
T ₄	168.56±3. 87	163.67±2. 31	158.67±2. 54	153.49±2. 81
T ₅	171.98±5. 23	163.22±4. 98	156.65±2. 78	150.23±3. 99
T ₆	167.43±3. 54	159.69±1. 56	151.39±3. 34	145.45±2. 32
T ₇	173.45±2. 41	166.32±4. 67	157.67±3. 89	152.06±4. 16
T ₈	170.56±3. 57	163.23±3. 16	154.89±4. 78	148.76±2. 45
T ₉	168.34±4. 45	159.34±4. 99	152.45±3. 87	144.93±3. 94
T ₀	167.45±3. 45	158.45±3. 74	144.5±2.5 6	124.50±3. 11

Table 3: Effect of different treatments on the fruit diameter (cm)

Treatments	0 Days	7 Days	14 days	21 days
T ₁	7.14±0.93	6.94±1.01	6.65±0.98	6.35±0.12
T ₂	7.10±0.75	6.91±1.21	6.71±0.75	6.44±1.02
T ₃	7.11±0.81	6.84±0.64	6.65±0.61	6.41±0.46
T ₄	7.54±0.83	7.33±0.76	7.10±1.45	6.86±0.76
T ₅	7.21±1.21	6.95±0.49	6.75±0.74	6.55±0.32
T ₆	7.19±0.92	6.87±0.40	6.57±0.98	6.25±0.15
T ₇	7.50±0.33	7.21±0.39	7.01±1.03	6.83±0.54
T ₈	7.01±0.41	6.74±0.86	6.45±0.21	6.20±0.33
T ₉	7.17±0.74	6.90±0.79	6.64±0.54	6.35±0.29
T ₀	7.24±0.75	6.74±1.22	6.21±0.66	5.65±1.11

Table 4: Effect of different treatments on the rind thickness of fruit

Treatments	0 Days	7 Days	14 days	21 days
T ₁	0.43±0.09	0.40±0.09	0.37±0.07	0.34±0.04
T ₂	0.39±0.03	0.36±0.04	0.32±0.04	0.29±0.03
T ₃	0.31±0.02	0.28±0.04	0.25±0.06	0.21±0.04
T ₄	0.42±0.08	0.41±0.09	0.39±0.04	0.35±0.06
T ₅	0.39±0.09	0.36±0.08	0.33±0.06	0.30±0.04
T ₆	0.43±0.02	0.40±0.07	0.35±0.04	0.31±0.05
T ₇	0.39±0.03	0.37±0.05	0.34±0.06	0.32±0.06
T ₈	0.43±0.04	0.41±0.07	0.34±0.06	0.30±0.05
T ₉	0.39±0.05	0.36±0.04	0.32±0.04	0.28±0.07
T ₀	0.41±0.06	0.35±0.06	0.31±0.05	0.25±0.06

Table 5: Effect of different treatments on juice content (%)

Treatment	0 Days	7 Days	14 days	21 days
T ₁	47.87±0.9 8	46.64±0.8 7	46.32±0.7 6	46.29±0.3 5
T ₂	48.18±0.8 7	47.76±0.8 6	47.43±0.8 7	47.15±0.8 7
T ₃	49.78±0.8 6	49.23±1.2 6	48.89±0.8 9	48.56±0.7 3
T ₄	52.67±0.6 7	51.98±1.5 6	51.54±0.8 3	51.31±0.8 1
T ₅	50.45±1.0 3	49.67±0.8 4	49.27±0.7 8	48.84±0.9 3
T ₆	51.65±0.4 5	50.34±0.6 7	49.75± .67	49.24±0.7 9
T ₇	49.56±0.9 3	48.76±0.8 7	48.43±0.6 6	48.29±0.8 8
T ₈	48.98±0.8 2	47.78±0.7 8	47.35±0.5 9	46.99±0.7 6
T ₉	49.54±0.8 7	48.96±0.9 7	48.54±1.3 8	48.12±0.4 5
T ₀	49.59±0.7 6	48.21±0.5 7	47.14±0.3 8	46.21±0.5 4

Table 6: Effect of different treatments on the colour of a fruit

Treatments	0 Days	7 Days	14 days	21 days
T ₁	9.57±0.45	8.59±0.37	6.24±0.67	5.44±0.32
T ₂	9.63±0.54	8.61±0.29	6.45±0.37	5.34±0.43
T ₃	9.55±0.49	8.63±0.43	6.52±0.43	5.67±0.43
T ₄	9.58±0.59	8.89±0.51	7.01±0.37	5.99±0.60
T ₅	9.70±0.23	8.78±0.87	6.92±0.58	5.81±0.46
T ₆	9.60±0.43	8.48±0.48	6.35±0.28	5.13±0.65
T ₇	9.50±0.66	8.54±0.32	6.49±0.79	5.38±0.77
T ₈	9.62±0.76	8.27±0.61	6.32±0.29	5.03±0.15
T ₉	9.71±0.49	8.10±0.58	6.01±0.88	4.98±0.25
T ₀	9.69±0.78	7.51±0.43	5.83±0.35	4.31±0.45

Table 7: Effect of different treatments on the flavor of fruit

Treatments	0 Days	7 Days	14 days	21 days
T ₁	9.40±0.35	8.59±0.57	6.39±0.86	5.02±0.64
T ₂	9.50±0.86	8.61±0.65	6.42±0.64	5.17±0.67
T ₃	9.60±0.87	8.63±0.88	6.82±0.68	5.26±0.79
T ₄	9.50±0.37	8.89±0.65	7.21±0.57	5.48±0.24
T ₅	9.70±0.74	8.78±0.43	7.92±0.32	5.41±0.63
T ₆	9.60±0.36	8.48±0.13	6.35±0.35	5.03±0.45
T ₇	9.50±0.83	8.94±0.15	6.49±0.46	5.19±0.87
T ₈	9.60±0.37	8.57±0.35	6.32±0.76	4.81±0.26
T ₉	9.70±0.64	8.30±0.65	6.01±0.64	4.98±0.65
T ₀	9.60±0.75	7.41±0.37	5.83±0.65	4.32±0.44

Table 8: Effect of different treatments on the overall acceptability of fruit

Treatments	0 Days	7 Days	14 days	21 days
T ₁	9.40±0.3	7.39±0.77	5.89±0.36	4.32±0.64
	5			
T ₂	9.50±0.8	7.51±0.65	5.82±0.94	4.57±0.67
	6			
T ₃	9.61±0.8	7.62±0.88	5.92±0.68	4.86±0.79
	7			
T ₄	9.50±0.4	7.89±0.65	5.97±0.57	5.18±0.24
	7			
T ₅	9.70±0.6	8.18±0.43	5.99±0.32	5.41±0.63
	4			

T ₆	9.60±0.5 6	6.41±0.73	4.36±0.35	3.23±0.95
T ₇	9.50±0.8 8	6.84±0.85	4.41±0.46	3.59±0.97
T ₈	9.60±0.3 9	5.97±0.35	4.22±0.76	3.11±0.96
T ₉	9.70±0.6 6	6.10±0.65	4.29±0.64	3.18±0.65
T ₀	9.60±0.6 7	5.41±0.37	3.73±0.67	2.12±0.94

Conclusion

Findings from the experiment showed that the perforated HDPE packaging content performed better than all of the other treatments. Further, it can be concluded that among the different treatments, the T₄ treatment proved superior as compare to other treatments in respect of maintaining physical and organoleptic traits. However, the minimum value was recorded for T₀ (Control) for most of the traits the traits studied. Therefore, treatment of T₄ (HDPE perforated) may be suggested as effective and economical treatment for improving shelf-life of mosambi fruits.

References

1. Bhullar JS, Farmahan HL, Agnihotri RP. Studies on storage behavior and extending the shelf life of Kinnow Mandarin. *Progressive Horticulture*. 1981;13(3-4):115- 119.
2. Deosthale YG, Pant KC. National Institute of Nutrition, Indian Council of Medical Research (ICMR); c2012. p. 47-58. "Food and Agriculture Organization" in India. <http://www.fao./home/en.com>
3. Indian database of horticulture. Horticultural Statistics at the Glance 2017. Government of India Ministry of Agriculture and Farmer's Welfare, Krishi Bhawan, New Delhi; c2017. p. 1-481.
4. Kaur S. Effect of high-density polyethylene, chemicals and curing on quality and storage of Kinnow. Thesis M.Sc. (Pomology). Punjab Agriculture University, Ludhiana, Punjab; c2000.
5. Nandigama, N. C. (2019). Energy-Efficient Virtual Machine Placement in Cloud Data Centers Using Reinforcement Learning-Enhanced Adaptive Greedy Dingo Optimization Algorithm. *Research Journal of Nanoscience and Engineering*, 3(1), 25–32. <https://doi.org/10.22259/2637-5591.0301006>.
6. Singh R. Effect of different packaging films and coatings on the shelf life and quality of Kinnow Fruit. Thesis M.Sc. (Pomology). Punjab Agricultural University, Ludhiana, Punjab, India; c2011.
7. Siddiqui S, Sharma RK, Gupta OP. Effect of Individual fruit wrapping by different materials on the shelf life of Guava Cv. Allahabad Safede. *Acta Horticulture*. 1997;735:535-545.

8. Nandigama, N. C. (2018). Deep Vision Networks for Multimodal Biometric Authentication: A Hybrid Feature-Level Fusion Approach with Machine Learning Optimization. *Research Journal of Nanoscience and Engineering*, 2(4), 22–29. <https://doi.org/10.22259/2637-5591.0204005>
9. Sonkar RK, Ladaniya MS. Individual film wrapping of Nagpur mandarin (*Citrus reticulata* Blanco) with heat- shrinkable and stretch-cling films for refrigerated Storage. *Journal of Food Science and Technology*. 1999;36(3):273-276.
10. Tatlı H, Özguven AI. The effect of some growth regulator treatments on the storage of Valencia. *Turkish Journal of Agriculture & Forestry*; c1999. p. 1033- 1042.
11. Tugwel BL, Chvyl WL. Modified Atmosphere packaging for citrus. Proceedings of the Australian post-harvest horticulture conference 'Science and technology for the fresh food revolution', Melbourne, Australia; c1996. p. 169-173.
12. Wills R, McGlasson B, Graham D, Joyce D. *Postharvest: An Introduction to the Physiology and Handling of Fruit, Vegetables, and Ornamentals* (2 edition); c2007.